



## INDONESIAN JOURNAL ON GEOSCIENCE

Geological Agency  
Ministry of Energy and Mineral Resources

Journal homepage: <http://ijog.geologi.esdm.go.id>  
ISSN 2355-9314, e-ISSN 2355-9306



### Geochemical Characterizations of Crude Oils in Pendopo High and Limau Graben, South Sumatra Basin

M. SYAIFUDIN<sup>1</sup>, and EDDY A. SUBROTO<sup>2</sup>

<sup>1</sup>Department Geology-FTM, Universitas Pembangunan Nasional “Veteran” Yogyakarta

<sup>2</sup>Department Geology-FITB, Institut Teknologi Bandung

Corresponding author: [msyaifudin@upnyk.ac.id](mailto:msyaifudin@upnyk.ac.id)

Manuscript received: November, 17, 2021; revised: November, 23, 2022;

approved: November, 13, 2023; available online: January, 29, 2024

**Abstract** – There are several oil fields in Pendopo High and Limau Graben, South Palembang Subbasin, South Sumatra Basin. Crude oils from the Pendopo High and Limau Graben have been analyzed using geochemical methods. Some or all of the following analyses were completed on twenty-four oils. Two analysis methods, namely the  $\delta^{13}\text{C}$  isotopic and biomarker compositions have been used to classify the oil types. Most of the oils are isotopically more depleted in  $\delta^{13}\text{C}_{\text{sat}}$  ( $> -28\%$ ) relative to the other four oil sample counterpart of the Pendopo High. All oil samples from The Pendopo High have Pr/Ph values  $< 4$  and those from the Limau Graben  $> 4$ . In addition, biomarker ratio for hopanes/steranes from the Limau Graben shows small values compared with the Pendopo High. Geochemical cross plots of probabilities can also be used to distinguish oils derived from different source inputs. In the Limau Graben, tricyclic terpane data shows terrestrial patterns, while the Pendopo High shows deltaic patterns. The Limau Graben is derived from humic kerogen, while the Pendopo High is more towards mixed kerogen (humic and sapropelic kerogen). The depositional condition in the Limau Graben is anoxic to suboxic to oxic and mostly oxic condition, predominated by higher plants. While Pendopo High condition is anoxic to suboxic, predominated by algae. Oil in the Pendopo High is interpreted to be deltaic oil, whereas oil in the Limau Graben is a typical fluvial.

**Keywords:** biomarker, deltaic oil, fluvial, South Sumatra Basin

© IJOG - 2024

#### How to cite this article:

Syaifudin, M. and Subroto, E.A., 2024. Geochemical Characterizations of Crude Oils in Pendopo High and Limau Graben, South Sumatra Basin. *Indonesian Journal on Geoscience*, 11 (1), p.1-13. DOI: [10.17014/ijog.11.1.1-13](https://doi.org/10.17014/ijog.11.1.1-13)

#### INTRODUCTION

Various geochemical methods have been applied to oil samples in attempts to provide information on the sources and geological conditions, including stable isotopes, gas chromatography, and gas chromatography-mass spectrometry (Robinson, 1987; Suseno *et al.*, 1992; ten Haven and Schiefelbein, 1995; Rashid *et al.*, 1998; Bishop, 2001; Peters *et al.*, 2005; Syaifudin *et al.*, 2015; and Syaifudin, 2016). In addition, the

literature contains various interpretations as to how isotopic compositions of oils can be related to the organic matter and depositional environments of their source rocks. For example, stable isotope compositions of crude oils and their saturate and aromatic hydrocarbon fractions have been used by petroleum geochemists to provide information on the depositional environment of the source rocks generating the oils, and the relationships among oils (Peters *et al.*, 2005 and references therein). Biomarker composition

can occur in different carbon ranges of crude oils (Peters *et al.*, 2005). In general, conventional biomarker compositions are characterized by sterane and triterpane families. Crude oil study, which utilizes the detailed geochemical analysis of a representative suite of samples, is an excellent way of identifying and comparing the samples that are sourced from the rock to a location of a relatively close area (Doust and Noble, 2008). Hence, their signatures are an invaluable tool for exploration because of their predictive capabilities.

Nederlandsche Koloniale Petroleum Maatschappij discovered Talang Akar field located in Pendopo High, South Sumatra Basin, Indonesia, in 1921 for producing the first oil from this area. The field was the biggest find in the country before World War-II. There are several oil fields have been discovered and most of them remain produce some oil to date. Geochemical study on oils from the Limau Graben and Pendopo High has been done by Suseno *et al.*, 1992. They found that the oils are characterized as fluvio-deltaic oil family, and were generated under some similarities in terms of geological phenomena (Bishop, 2001). This study was done to confirm that there are differences between crude oil samples in the Limau Graben and Pendopo High. The crude oil samples contained in the Limau Graben seem to have been deposited relatively more towards an oxidation condition ( $Pr/Ph > 4$ ) as well as have higher marker than those in the Pendopo High (Tables 1 and 2). In the Limau Graben, tricyclic terpane data shows terrestrial patterns ( $C_{19}$  and  $C_{20}$  tricyclic terpanes are relatively abundant), while those in the Pendopo High indicate delta patterns ( $C_{19}$ ,  $C_{20}$ , and  $C_{23}$  tricyclic terpanes are abundant) (Tables 3 and 4).

This paper focuses to discuss the successfully combined use of isotopic and biomarker indicators to distinguish and to separate the Pendopo High and Limau Graben crude oils based on their closed geographic locations. Although this preliminary work is based on a limited sample suite, the useful results justify an expanded study using more oils and analytical parameters.

The studied area is located in the Pendopo High area and Limau Graben, South Palembang Subbasin, South Sumatra Basin (Figure 1).

## METHODS AND MATERIALS

The analytical methods used in this study were capillary gas chromatography (GC); a standard gravitational column chromatography for hydrocarbon fractionation and saturate GC-MS for biomarker determination.

The whole oil sample was previously analyzed using a fast GC to find alkanes distribution. Fingerprinting analysis of the whole oil was carried out by using an Agilent 6890 N GC instrument coupled to a flame ionization detector (FID). The GC was fitted with a 10 m x 0.21mm i.d. DB-1 (J and W) fused-silica capillary column. Samples were injected using an Agilent 7673 auto sampler, with split/splitless mode injector.

For GC-MS analysis, the oil samples were separated into saturated hydrocarbon, aromatic hydrocarbon, and nonhydrocarbon fraction (polar compounds) by a preparative column chromatography. Isolation of branched and cyclic fraction from saturated hydrocarbons was carried out using a packed activated silicalite chromatographic column. Then the branched and cyclic fractions were introduced to a GC-MS instrument.

GC-MS analyses of saturate fraction were performed using an Agilent 6890 Gas Chromatograph (GC) coupled to an Agilent 5973 series Mass Selective Detector (MSD) - computer data system (Chemstation). The GC was fitted with a 60 m x 0.25 mm i.d. DB-5MS (J and W) fused-silica capillary column. Samples were injected using an Agilent 7673 auto sampler, with split/splitless mode injector. The MS condition was ionization mode: electron impact (EI), EM voltage was 1980 Volt; electron energy was 70 eV and source temperature 250°C.

Examination of fingerprint signatures for all samples were based on pristane, phytane of the whole oil chromatograms and triterpanes ( $m/z$  191), and steranes ( $m/z$  217) of saturated fractions.

Geochemical Characterizations of Crude Oils in Pendopo High and Limau Graben, South Sumatra Basin  
(M. Syaifudin and E.A. Subroto)

Table 1. Biomarker Data of Oils in the Pendopo High

Wells in Pendopo High	$\delta^{13}\text{C}$ sats	$\delta^{13}\text{C}$ aroms	Ph/nC <sub>18</sub>	Pr/nC <sub>17</sub>	Pr/Ph	Tot Hop/Tot Ster
Pandawa-1	-27,36	-26,63	0,23	0,68	3,10	92,90
Pandawa-1	-29,50	-27,33	0,19	0,57	3,26	0
Pandawa-1	-29,11	-27,12	0,20	0,56	3,06	19,36
Pandawa-2	-26,40	-26,20	0,21	0,72	3,28	12,75
Pandawa-3	-28,70	-27,70	0,32	0,89	2,84	9,82
Pandawa-4	-27,10	-27,20	0,31	1,08	3,74	7,29
Pandawa-5	0	0	0,34	1,09	3,38	7,56
Pandawa-6	-27,90	-27,60	0,26	1,32	3,95	9,46

Table 2. Biomarker Data of Oils in the Limau Graben

No	Well in Limau Graben	$\delta^{13}\text{C}$ sats	$\delta^{13}\text{C}$ aroms	Ph/nC <sub>18</sub>	Pr/nC <sub>17</sub>	Pr/Ph	Tot Hop/Tot Ster
9	Orange-1	-28,7	-27,20	0,65	2,89	4,02	3,88
10	Orange-2	-29,50	-27,20	0,75	7,31	10,56	3,94
11	Orange-3	0	0	0,41	1,81	5,22	4,31
12	Orange-4	-29,80	-27,80	0,15	1,04	7,75	2,84
13	Orange-4	-29,30	-28,10	0,13	1,12	9,33	3,30
14	Orange-4	-29,70	-28,00	0,14	1,17	7,58	3,03
15	Orange-5	-28,30	-27,00	0,60	4,94	14,76	33,3
16	Orange-5	-29,10	-27,70	0,23	2,07	11,44	9,10
17	Orange-6	-29,00	-27,30	0,48	6,73	4,21	3,78
18	Orange-6	-29,90	-28,30	0,21	2,30	4,05	4,29
19	Orange-7	-29,40	-29,90	0,34	1,76	4,96	2,84
20	Orange-8	-29,10	-27,40	0,48	1,55	4,01	3,58
21	Orange-9	-28,90	-27,10	0,31	1,87	6,50	2,33
22	Orange-10	-29,30	-27,30	0,38	1,33	7,90	5,90
23	Orange-11	0	0	0,62	4,83	6,50	2,33
24	Orange-12	0	0	0,31	2,24	7,90	5,90

Table 3. Tricyclic Terpane Data of Oils in the Pendopo High

Wells	Norm% Tricyclics Terpane						
	B	C	D	E	F	G	H
	C19	C20	C21	C22	C23	C24	C25
Pandawa-2	18	16	21	0	22	16	7
Pandawa-3	20	161	17	0	20	18	10
Pandawa-4	18	15	17	3	18	18	12
Pandawa-5	18	15	16	0	19	19	12
Pandawa-6	19	15	14	5	17	17	14

Table 4. Tricyclic Terpane Data of Oils in the Limau Graben

Wells	Norm% Tricyclics Terpane						
	B	C	D	E	F	G	H
	C19	C20	C21	C22	C23	C24	C25
Pandawa-2	18	16	21	0	22	16	7
Pandawa-3	20	161	17	0	20	18	10
Pandawa-4	18	15	17	3	18	18	12
Pandawa-5	18	15	16	0	19	19	12
Pandawa-6	19	15	14	5	17	17	14




Figure 1. Location maps of (a) South Sumatra Basin and (b) studied area.

In this study, the oil analysed consist of twenty samples of carbon isotopes and twenty-four samples of biomarkers. Characterization has been based on qualitative and quantitative data. Qualitative data comprised evaluations which were based on chromatograms and mass-fragmentograms. Whereas the quantitative data consists of a series of cross plots, eg. Cross plot of carbon isotope  $\delta^{13}\text{C}$  saturates-aromatics, distribution of  $\text{C}_{27}\text{-C}_{28}\text{-C}_{29}$  sterane,  $\text{Pr/nC}_{17}\text{-Ph/nC}_{18}$ ,  $\text{Pr/Ph-Pr/nC}_{17}$ , carbon isotope  $\delta^{13}\text{C}$  saturates- $\text{Pr/Ph}$ ,  $\text{Pr/Ph-total hopane/total sterane}$ , and ratio of  $\text{C}_{26}/\text{C}_{25}$  (tricyclic).

### Regional Structural Geology

Geological structures that control the regional South Sumatra were influenced by three tectonic phases (Pulunggono *et al.*, 1992):

- Compression (Upper Jurassic - Lower Cretaceous)
- Tension (Upper Cretaceous - Lower Tertiary)
- Compression (Middle Miocene - Recent) .

The first phase: Started in Upper Jurassic - Lower Cretaceous, characterized by the subduction of India-Australia Plate as a movement

mechanism to yield primary stress to the Sundaland trending N 30° W. This subduction resulted simple shear (N 300° E) as a strike-slip fault that was actively moved laterally. This was assumed as the cause of linearity trending N-S as an antithetic fault which was inactive.

The second phase: Commenced during Upper Cretaceous-Lower Tertiary, characterized by the change of subduction trend of the India-Australia Plate into N-S. This event resulted in the formation of some geological structures (fractures) caused by a tension force as a linearity with N-S direction. This phenomenon caused the formation of grabens and depressions, such as Benakat Gully. Initiation of graben filling with Tertiary sediments was started.

The third phase: Commenced in the Middle Miocene-present, shown with, again, the change of the subduction direction into N 6° E, causing rejuvenation and inversion processes on the paleostructures by Plio-Pleistocene (N 330° E) and the uplifting of Barisan Mountains and also the formation of some thrust faults with Lematang Fault pattern.



## Regional Stratigraphy

Based on the tectonostratigraphy framework, Ryacudu (2008) divides Early Tertiary rock units in South Sumatra Basin as follows (Figure 2):

- Pre-rift sequences

This sequence consists of volcanic rocks of Kikim Formation and pre-Tertiary rocks. Kikim Formation are the oldest Tertiary rocks in South Sumatra Basin, consisting of volcanic rocks such as volcanic breccia, agglomerate, andesitic tuffs, and igneous rocks (as intrusions and lava flows). The age of Kikim Formation based on dating K-Ar is 54-30 Ma (Paleocene-Early Oligocene, Ryacudu, 2008).

- Syn-rift sequences

Syn-rift sequence consists of Lahat Group comprising Lemat and Benakat Formations with interfingering relationship (Ryacudu, 2008). The main constituent of Lemat Formation is coarse clastic rocks (sandstone) with Tuff and Conglomerate Members, while Benakat Formation is dominated by fine clastic rocks (shale). The Lemat Formation deposited in a fluvial environment. However, in the Pendopo High, Lemat Formation is likely to have a fluvio-deltaic environment.

Shale of Benakat Formations interpreted as lake (lacustrine) sediments.

- Post-rift sequences

Tanjungbaru Formation, originally considered as a GRM (Gritsand Member) formerly known as a member of Talangakar Formation. This unit is dominated by conglomeratic sandstone deposition system as the result of a braided river. Notwithstanding, in Pendopo High, Lower Talangakar Formation (GRM) is likely to have fluvio-deltaic environment. This member of Talangakar Formation commonly referred as a TRM (Transition Member) is proposed as Talangakar Formation. This formation consists of alternating sandstones and shales, with thin coal interbedded, deposited in a transition environment, *i.e.* a delta system to a shallow marine of Early Miocene. Baturaja Formation, Early Miocene (N5-N6) in age, composed of bioclastic limestone, calcarenite, bioclastic sandy limestones, and reefal bioherm with interbedded of calcareous shale, was deposited on the a carbonate platform. Upwards, Early Miocene to Middle Miocene Gumai Formation, is composed of calcareous mudstone containing fossil planktonic *Globigerina* and shales with




Figure 2. Regional stratigraphy of South Sumatra Basin (modified from Ryacudu, 2008).

glaucconitic quartz sandstones. Deposited conformity on top of the Gumai Formation is Palembang Group, comprising of Air Benakat, Muara Enim, and Kasai Formations. Furthermore, the marine condition was getting shallow and then the overlying Kasai Formation was deposited in a terrestrial environment.

## RESULTS AND DISCUSSION

### Characterization of Oils in the Pendopo High and Limau Graben Areas

A qualitative comparison of biomarker characterizations between crude oil samples in the Pendopo High and Limau Graben is presented in Figure 3.

Based on data of  $H_{29}$  and  $H_{30}$  (hopane) distribution, the pattern of  $H_{29} < H_{30}$  indicates clastic sediments, while  $H_{29} > H_{30}$  is evaporates-carbonate sediment (Zumberge, 1984; Connan *et al.*, 1986; Price *et al.*, 1987; Waples and Machihara, 1991; Peters and Moldowan, 1993; Peters *et al.*, 2005; Hakimi *et al.*, 2011; Grosjean *et al.*, 2012; Xiangchun

*et al.* 2013; ten Haven *et al.*, 1988 vide Rabbani *et al.*, 2014; Syaifudin *et al.*, 2015; Wang *et al.*, 2015; and Syaifudin, 2016). The crude oil samples from Pendopo High and Limau Graben are clastic sediments since it shows as pattern of  $29_H < 30_H$ .

From data of homohopane distribution which decreased regularly from  $C_{31}$  to  $C_{35}$ , the crude oil samples from Pendopo High and Limau Graben are interpreted as depositional environment which associated with clastic sediments (Waples and Machihara, 1991) or more oxidizing conditions (Peters and Moldowan, 1993; Hakimi *et al.*, 2011).

A terrestrial environment is characterized by high  $C_{19}$  and  $C_{20}$  tricyclic terpane (Philp and Gilbert, 1986; Peters and Moldowan, 1993; Hanson *et al.*, 2000; Grice *et al.*, 2001; George *et al.*, 2004; Volk *et al.*, 2005; Zhang and Huang 2005; Peters *et al.*, 2005; George *et al.*, 2007; Hao *et al.*, 2011; Asif *et al.*, 2011; Adedosu *et al.*, 2012; Alberdi *et al.*, 2001 vide Rabbani *et al.*, 2014; Tao *et al.*, 2015; Wang *et al.*, 2015). Crude oil samples from the Limau Graben show  $C_{19}$  and  $C_{20}$  tricyclic terpanes are more abundant (Figure




Figure 3. Comparison of biomarker characterization qualitatively between crude oils in the Pendopo High (left) and Limau Graben (right).

4).  $C_{23}$  tricyclic terpene is often predominant in crude oils of a marine source (Aquino Neto *et al.*, 1983; Zumberge, 1987; Burwood *et al.*, 1992; Peters and Moldowan, 1993; Hanson *et al.*, 2000; Zhang and Huang, 2005; Hakimi *et al.*, 2011; Tao *et al.*, 2015). Crude oil samples in the Pendopo High besides show predominant  $C_{19}$  and  $C_{20}$  tricyclic terpanes, they also contain relatively high concentration of  $C_{23}$  tricyclic terpene.

Based on these data, crude oil in the Limau Graben is interpreted to be originated from terrestrial source rocks, and the crude oil samples from the Pendopo High are interpreted to be derived from deltaic source rocks (Figure 4).

### Quantitative Characterization

Pristane/phytane (Pr/Ph) ratios have been used to assess the redox potential of the depositional environment and the source of organic matter (Didyk *et al.*, 1978; Tissot and Welte, 1984; Zhang and Huang, 2005; Peters *et al.* 2005; Duan *et al.*, 2008; Hao *et al.*, 2011; Hakimi *et al.*, 2011; Adedosu *et al.* 2012; Cheng *et al.*, 2013; and Tao *et al.*, 2015) or reflect the relationship between contributing organisms and the chemistry of the environment (Mello and Maxwell, 1990).

The cross plot of Pr/Ph - Pr/nC<sub>17</sub>, as proposed by Robinson (1991) (Figure 5a), shows that crude oil samples from Pendopo High fall into an anoxic




Figure 4. Comparison of tricyclic terpanes between crude oils in the Pendopo High (left) and the Limau Graben (right).



to suboxic category and was dominated by algae. While crude oil samples from Limau Graben fall into anoxic to oxic, mostly oxic, predominated by higher plants. The crude oil samples contained in the Limau Graben seem to have been deposited relatively more towards an oxidation condition ( $Pr/Ph > 4$ ) as well as to have the higher marker than the crude oil samples in the Pendopo High.

The cross plot  $Pr/nC_{17} - Ph/nC_{18}$  (see Figure 5b) as proposed by Connan and Cassou (1980), shows that crude oil samples from Pendopo High and Limau Graben derived from humic kerogen, but in the Pendopo High they are more towards mixed kerogen (humic and sapropelic kerogen) (Sabra, 2021). The cross plot of carbon isotope  $\delta^{13}C$  saturates - aromatics as proposed by Sofer (1984), shows that crude oil samples from Pendopo High (Figure 6a) are mostly derived from alga, and the saturate fraction are heavier than crude oil samples from Limau Graben.

The cross plot of carbon isotope  $\delta^{13}C$  saturates -  $Pr/Ph$ , as proposed by Bishop (2001) (Figure 6b), shows that crude oil samples from Limau Graben are more oxic than crude oil samples from Pendopo High.




Figure 5. Cross plot of  $Pr/Ph - Pr/nC_{17}$  (a),  $Pr/nC_{17} - Ph/nC_{18}$  (b). Crude oils in the Pendopo High and Limau Graben.




Figure 6. Cross plot of (a) carbon isotope  $\delta^{13}C$  saturates - aromatics and (b) carbon isotope  $\delta^{13}C$  saturates -  $Pr/Ph$ , crude oils in the Pendopo High and Limau Graben.

The cross plot of  $Pr/Ph$  - total hopane/sterane crude oils in the Pendopo High and Limau Graben as proposed by Robinson (1991) (Figure 7), shows that crude oil samples from Pendopo High fit to an anoxic to suboxic terrestrial influence, while crude oil samples from Limau Graben mostly fall into a highly oxidizing terrestrial.




Figure 7. Cross plot of  $Pr/Ph$  - total hopane/sterane, crude oils in the Pendopo High, and Limau Graben.

The isopach map of Lemat and Lower Talangarakar Formations (Figure 8) as source rocks, and the possibility of oil migration pathways prior Plio-Pleistocene, while the Top structure map of Talangarakar Formation, and the possibility of oil






Figure 8. Isopach map of the Lemat and Lower TAF as source rocks, and the possibility of oil migration pathways prior Plio-Pleistocene.

migration pathways after Plio-Pleistocene are exhibited in Figure 9. Based on these data, crude oil samples from the Pendopo High are interpreted to be a deltaic oil




Figure 9. Top structure map of the Talangakar Formation, and the possibility of oil migration pathways after Plio-Pleistocene.

family, whereas crude oil samples from the Limau Graben are categorized as the fluvial oil family.

### CONCLUSIONS

Based on tricyclic terpane data, the crude oil samples from Pendopo High are interpreted to be derived from a deltaic environment, while those from Limau Graben are deciphered derived from a fluvial environment.

The crude oil samples from Limau Graben seem to have been deposited relatively more towards an oxidation condition ( $\text{Pr/Ph} > 4$ ) as well as have higher plant marker than the crude oil samples from Pendopo High.

Crude oil samples from Limau Graben and Pendopo High have been derived from a humic kerogen, however those found in the Pendopo High indicate more towards mixed kerogen (humic and sapropelic kerogen).

The  $\delta^{13}\text{C}$  saturate fraction from crude oil samples from Pendopo High is heavier than crude oil samples from Limau Graben ( $\delta^{13}\text{C}$  saturate fraction Pendopo High -26.4 to -29.5, while in Limau Graben -28.3 to -29.8).

In general, crude oil samples from the Pendopo High are interpreted to be deltaic oil family, whereas crude oil samples from Limau Graben are typical a fluvial oil family.

### ACKNOWLEDGEMENTS

The authors would like to thank the management of the Directorate General of Oil and Gas and Medco EP for approving to publish this paper.

### REFERENCES

- Adedosu, T.A., Sonibare, O.O., Tuo, J., and Ekundayo, O., 2012. Biomarkers, carbon isotopic composition and source rock potentials of Awgu coals, middle Benue trough, Nigeria. *Journal of African Earth Sciences*, 66-67. p.13-21. DOI: 10.1016/j.jafrearsci.2012.03.006.
- Aquino Neto, F.R., Trendel, J.M., Restle, A., Connan, J., and Albrecht, P.A., 1983. Occurrence and formation of tricyclic and tetracyclic terpanes in sediments and petroleum. *In: Bjarøy, M., Albrecht, P., Conford, C., de Groot, K., Eglinton, G., Galimov, E., Leythaeuser, D., Pelet, R., Rullkoter, J., and Speers, G. (eds.), Journal Advances in Organic Geochemistry*, John Wiley, Chichester, p.659-667.
- Asif, M., Fazeelat, T., and Grice, K., 2011. Petroleum geochemistry of the Potwar Basin, Pakistan: 1. Oil-oil correlation using biomarkers,  $\delta^{13}\text{C}$  and  $\delta\text{D}$ . *Journal Organic Geochemistry*, 42, p.1226-1240. DOI: 10.1016/j.orggeochem.2011.08.003.
- Bishop, M.G., 2001. South Sumatra Basin Province, Indonesia: The Lahat/Talang Akar-Cenozoic Total Petroleum System. *U.S. Geological Survey, Open-File Report 99-50-s*. DOI: 10.3133/ofr9950S
- Burwood, R., Leplat, P., Mycke, B., and Paulet, J., 1992. Rifted margin source rock deposition: a carbon isotope and biomarker study of a west African Lower Cretaceous "lacustrine" section. *Journal Organic Geochemistry*, 19 (1-3), p.41-52. DOI: 10.1016/0146-6380(92)90026-T
- Cheng, P., Xiao X.M., Tian H., Huang B.J., Wilkins R.W.T., and Zhang, Y.Z., 2013. Source controls on geochemical characteristics of crude oils from the Qionghai Uplift in the western Pearl River Mouth Basin, offshore South China Sea. *Journal Marine and Petroleum Geology*, 40, p.85-98. DOI: 10.1016/j.marpetgeo.2012.10.003.
- Connan, J. and Cassou, A.M., 1980. Properties of gases and petroleum liquids derived from terrestrial kerogen at various maturation levels. *Journal Geochimica et Cosmochimica Acta*, 44 (1), p.1-23. DOI: 10.1016/0016-7037(80)90173-8
- Connan, J., Bouroullec, J., Dessort, D., and Albrecht, P., 1986. The microbial input in carbonate-anhydrite facies of a sabkha palaeoenvironment from Guatemala: a molecular

- approach. *Journal Organic Geochemistry*, 10 (1-3), p.29-50. DOI: 10.1016/0146-6380(86)90007-0
- Didyk, B.M., Simoneit, B.R.T., Brassell, S.C., and Eglinton, G., 1978. Organic geochemical indicators of palaeoenvironmental conditions of sedimentation. *Journal Nature*, 272, p.216-222. DOI: 10.1038/272216a0
- Doust, H. and Noble R.A., 2008. Petroleum systems of Indonesia, *Journal Marine and Petroleum Geology*, 25, p.103-129. DOI: 10.1016/j.marpetgeo.2007.05.007.
- Duan, Y., Wang, C.Y., Zheng, C.Y., Wu, B.X., and Zheng, G.D., 2008. Geochemical study of crude oils from the Xifeng oilfield of the Ordos basin, China. *Journal of Asian Earth Sciences*, 31, p.341-356. DOI: 10.1016/j.jseaes.2007.05.003.
- George, S.C., Lisk, M., and Eadington, P.J., 2004. Fluid inclusion evidence for an early, marine-sourced oil charge prior to gas-condensate migration, Bayu-1, Timor Sea, Australia. *Journal Marine and Petroleum Geology*, 21 (9), p.1107-1128. DOI: 10.1016/j.marpetgeo.2004.07.001
- George, S.C., Volk, H., Ahmed, M., Pickel, W., Allan, T., 2007. Biomarker evidence for two sources for solid bitumens in the Subu wells: Implications for the petroleum prospectivity of the East Papuan Basin, *Journal Organic Geochemistry*, 38 (4), p.609-642. DOI: 10.1016/j.orggeochem.2006.12.002
- Grice, K., Audino, M., Alexander, R., Boreham, C.J., Kagi, R.I., 2001. Distributions and stable carbon isotopic compositions of biomarkers in torbanites from different palaeogeographical locations. *Journal Organic Geochemistry*, 32 (10), p.195-211. DOI: 10.1016/S0146-6380(01)00087-0
- Grosjean, E., Love G.D., Kelly A.E., Taylor, P.N., and Summons R.E., 2012. Geochemical evidence for an Early Cambrian origin of the 'Q' oils and some condensates from north Oman, *Journal Organic Geochemistry*, 45, p.77-90. DOI: 10.1016/j.orggeochem.2011.12.006
- Hakimi, M.H., Abdullah, W.H., and Shalaby, M.R., 2011. Organic geochemical characteristics of crude oils from the Masila Basin, eastern Yemen. *Journal Organic Geochemistry*, 42 (5), p.465-476. DOI: 10.1016/j.orggeochem.2011.03.015.
- Hanson, A.D., Zhang, C., Moldowan, J.M., Liang, D.G., and Zhang, B.M., 2000. Molecular organic geochemistry of the Tarim Basin, north-west China. *American Association of Petroleum Geologists Bulletin*, 84, p.1109-1128.
- Hao, F., Zhou, X., Zhu, Y., and Yang, Y., 2011. Lacustrine source rock deposition in response to co-evolution of environments and organisms controlled by tectonic subsidence and climate, Bohai Bay Basin, China. *Journal Organic Geochemistry*, 42, p.323-339. DOI: 10.1016/j.orggeochem.2011.01.010.
- Mello, M.R. and Maxwell, J.R., 1990. Organic geochemical and biological marker characterization of source rocks and oils derived from Lacustrine environments in the Brazilian continental margin, in Lacustrine basin exploration, Case studies and modern analogs. In: Katz, B.J. (ed.), *AAPG Memoir*, 50, p.77-97. DOI: 10.1306/M50523C5
- Peters, K.E. and Moldowan J.M., 1993. *The Biomarker guide, Interpreting molecular fossils in petroleum and ancient sediments*. Prentice Hall, Englewood Cliffs, New Jersey, p.1-363.
- Peters, K.E., Walters, C.C., and Moldowan J.M., 2005. *The Biomarker guide*, (1, 2), Cambridge University press, 1155pp.
- Philp, R.P. and Gilbert, T.D., 1986. Biomarker distributions in oils predominantly derived from terrigenous source material. In: Leythaeuser, D. and Rullkötter, J. (eds.), *Advances in Organic Geochemistry 1985. Journal Organic Geochemistry*, 10 (1-3), p.73-84. DOI: 10.1016/0146-6380(86)90010-0
- Price, P.L., Sullivan, T.O., and Alexander, R., 1987. The Nature and occurrence of oil in Seram, Indonesia. *Proceedings of Indonesia Petroleum Association, 16<sup>th</sup> Annual Convention, Jakarta, Indonesia*.
- Pulunggono, A., Agus, H. S., and Kosuma, C. G., 1992. Pre-Tertiary and Tertiary fault systems as a framework of the South Sumatra basin;

- A Study of SAR maps. *Proceedings Indonesian Petroleum Association, 21<sup>st</sup> Annual Convention & Exhibition, Jakarta, Indonesia*, p.339-360.
- Rabbani, A.R., Kotarba, M.J., Baniasad, A.R., Hosseiny, E., Wieclaw, D., 2014. Geochemical characteristics and genetic types of the crude oils from the Iranian sector of the Persian Gulf. *Journal Organic Geochemistry*, 70, p.29-43. DOI: 10.1016/j.orggeochem.2014.02.010
- Rashid, H., Sosrowidjojo, I.B. dan Widiarto, F.X., 1998. Musi platform and Palembang high: A New look at the petroleum system. *Proceedings of Indonesian Petroleum Association, 26<sup>th</sup> Annual Convention*, Jakarta, Indonesia, 265-276.
- Robinson, K.M. 1987. An Overview of source rocks and oils in Indonesia. *Proceedings of Indonesian Petroleum Association, 16<sup>th</sup> Annual Convention*, Jakarta, Indonesia, p.97-122.
- Robinson, 1991. Geochemistry of oil and gas samples from Gegas 1, South Sumatra PSC, Indonesia, PT Geoservices (Ltd.), Report No 2007/0606/LAB, prepared for BPMIGAS MEDCO E&P. Unpublished.
- Ryacudu, R., 2008. Tinjauan stratigrafi Paleogen Cekungan Sumatra Selatan, Sumatra. *Stratigraphy Workshop IAGI*, p.99-114.
- Sabra, E., 2021. Evaluation of Hydrocarbon source rock potensial and organic geochemistry in North Arafura Shelf, Papua (Indonesia). *Indonesian Journal on geoscience* 8 (3), p. 401-416. DOI: 10.17014/ijog.8.3.401-416
- Sofer, Z., 1984. Stable carbon isotope compositions of crude oils: application to source depositional environments and petroleum alteration. *American Association of Petroleum Geologists, Bulletin*, 68, p.31-49.
- Suseno, P.H., Zakaria, Mujahidin, N., and Subroto, E.A., 1992. Contribution of Lahat formation as hydrocarbon source rock in South Palembang area, South Sumatra, Indonesia. *Proceedings of Indonesian Petroleum Association, 21<sup>st</sup> Annual Convention*, Jakarta, Indonesia, p.325-337.
- Syaifudin, M., 2016. Organic geochemical characteristic of crude oils from Limau Graben, South Sumatra Basin. *Journal of Geoscience, Engineering, Environment, and Technology*, 1 (1), p.27-34. DOI: 10.24273/jgeet.2016.11.3.
- Syaifudin, M., Subroto, E.A., Noeradi, D., and Kesumajana, A.H.P., 2015. Characterization and correlation study of source rocks and oils in Kuang area, South Sumatra Basin: the potential of Lemat Formation as hydrocarbon source rocks. *Proceedings of Indonesian Petroleum Association, 39<sup>th</sup> Annual Convention & Exhibition*, Jakarta, Indonesia.
- Tao, S., Wang, C., Du, J., Liu, L., and Chen, Z., 2015. Geochemical application of tricyclic and tetracyclic terpane biomarkers in crude oils of NW China. *Journal Marine and Petroleum Geology*, 67, p.460-467. DOI: 10.1016/j.marpetgeo.2015.05.030.
- ten Haven and Schiefelbein, 1995. The petroleum systems of Indonesia. *Proceedings of Indonesian Petroleum Association, 24<sup>th</sup> Annual Convention*, Jakarta, Indonesia, p.443-459.
- Tissot, B.P. and Welte, D.H., 1984. *Petroleum formation and occurrence, Second revised and enlarge edition*, Springer-Verlag, Berlin-Heidelberg-New York-Tokyo, 699pp.
- Volk, H., George, S., Middleton, H., and Schofield, S., 2005. Geochemical comparison of fluid inclusion and present-day oil accumulations in the Papuan Foreland-evidence for previously unrecognised petroleum source rocks. *Journal Organic Geochemistry*, 36, p.29-51. Elsevier
- Wang, Q., Hao, F., Xu, C., Wang, Y., and Zou, H., 2015. Geochemical characterization of QHD29 oils on the eastern margin of Shijiutuo uplift, Bohai Sea, China: Insights from biomarker and stable carbon isotope analysis. *Journal Marine and Petroleum Geology*, 64, p.266-275. DOI: 10.1016/j.marpetgeo.2015.03.010.
- Waples, D.W. and Machihara, T., 1991. Biomarkers for geologist - A Practical guide to the application of steranes and triterpanes in petroleum geology. *AAPG Methods in Exploration*, 9, The American Association of Petroleum Geologist, 91pp.



- Xiangchun, C., Wang, T.G., Qiming, L., Bin, C., and Xiaowan, T., 2013. Geochemistry and possible origin of petroleum in Palaeozoic reservoirs from Halahatang depression. *Journal of Asian Earth Sciences*, 74, p.129-141. DOI: 10.1016/j.jseaes.2013.03.024.
- Zhang, S. and Huang, H., 2005. Geochemistry of Palaeozoic marine petroleum from the Tarim Basin, NW China: Part 1 Oil family classification. *Journal Organic Geochemistry*, 36, p.1204-1214. DOI: 10.1016/j.orggeochem.2005.01.013.
- Zumberge, J.F., 1984. Source rocks of the La Luna Formation (Upper Cretaceous) in the Middle Magdalena Valley, Colombia. In: Palacas, J.G. (ed.), *Geochemistry and Source Rock Potential of Carbonate Rocks. American Association of Petroleum Geologist Studies in Geology #18*, Tulsa, p.127-133.
- Zumberge, J.F., 1987. Terpenoid biomarker distributions in low maturity crude oils. *Journal Organic Geochemistry*, 11, Elsevier, p.479-496.