



INDONESIAN JOURNAL ON GEOSCIENCE

Geological Agency
Ministry of Energy and Mineral Resources

Journal homepage: <https://ijog.geologi.esdm.go.id>
ISSN 2355-9314, e-ISSN 2355-9306



Combining Lithium Isotope Systematics, Stable Isotopes, and Major Elements as a Powerful Tool for Detection of the Slab-Dehydrated Fluids in Southwestern Japan

^{1,2}ROFIQUL UMAM

¹Faculty of Life and Environmental Sciences, University of Tsukuba, Japan

²Center for Research in Radiation, Isotopes, and Earth System Sciences (CRiES),
University of Tsukuba, Japan

Corresponding author: rofiqul.umam@ied.tsukuba.ac.jp

Manuscript received: October 15, 2024; revised: May 18, 2025;
approved: May, 19, 2026; available online: August, 4, 2026

Abstract – This study aims to evaluate whether an integrated geochemical approach using lithium isotope systematics ($\delta^7\text{Li}$), Li and Cl concentrations, Cl/Li ratios, stable isotopes (δD and $\delta^{18}\text{O}$), and major-element compositions can provide useful constraints on the contribution of thermal water to the hydrothermal system. The study is based entirely on previously published geochemical data. Lithium isotope ($\delta^7\text{Li}$) data for four Arima samples reported by Umam *et al.* (2022) were re-evaluated together with Li and Cl concentration data reported by Umam *et al.* (2022) and Kusuda *et al.* (2014). Major-element concentrations and stable isotope (δD and $\delta^{18}\text{O}$) data for Arima hot spring waters reported by Kusuda *et al.* (2014) were also integrated into the analysis. The four Arima samples with available $\delta^7\text{Li}$ data have $\delta^7\text{Li}$ values $< 5\text{‰}$ and Cl/Li ratios < 1000 and are interpreted as representing thermal water. Although $\delta^7\text{Li}$ data are not available for the Arima samples reported by Kusuda *et al.* (2014), 11 of these samples satisfy the proposed thermal-water criteria of Li concentrations $> 0.1\text{ mg/L}$, Cl concentrations $> 1000\text{ mg/L}$, and Cl/Li ratios < 1000 . One sample is interpreted as representing mixing water and has a Cl concentration of 435 mg/L . These results suggest that integrating lithium isotope systematics with Li and Cl concentrations, Cl/Li ratios, stable isotopes, and major-element compositions may provide a useful geochemical framework for evaluating the characteristics and possible contribution of slab-dehydrated fluids to the hydrothermal system.

Keywords: Arima hot spring waters, slab-dehydrated fluids, lithium isotope systematics, stable isotopes, major elements, powerful geochemical analysis

© IJOG - 2026

How to cite this article:

Umam, R., 2026. Combining Lithium Isotope Systematics, Isotope Stable, and Major Elements as a Powerful Tool for Detection of The Slab-Dehydrated Fluids in Southwestern Japan. *Indonesian Journal on Geoscience*, 13 (3), p.393-407. DOI: [10.17014/ijog.13.3.393-407](https://doi.org/10.17014/ijog.13.3.393-407)

INTRODUCTION

Background

For more than a decade, the emergence of fluids originating from the Earth's mantle, especially in subduction zone areas (slab-dehydrated fluids/thermal waters), has been considered to play an important role in geodynamic processes (Nakajima and Hasegawa, 2007; Shiomi and Park,

2008). Its role in the process of island origin has been known for a long time. In addition, the activity of the emergence of slab-dehydrated fluids has a dramatic effect on various rock properties and indicates the control of other geophysical processes in the area around the slab (Morikawa *et al.*, 2008). Specifically, slab-dehydrated fluids are fluids that are formed from the process of slabs entering the Earth, are dehydrated, and undergo

Indexed by: SCOPUS

a long process of water-rock interaction before finally being released through faults or fractures around subduction zones (Morikawa *et al.*, 2008; Stern, 2018; Umam *et al.*, 2022).

In the process, hydrated areas differ from dehydrated areas, such that the low-velocity zone beneath the lithosphere, which was originally interpreted as a partial melting zone, can be viewed as a hydrated zone (Wyllie, 1974). Meanwhile, zones with higher temperatures, pressure, and trapping can be considered dehydration zones (Harlaux *et al.*, 2017; Tanimizu *et al.*, 2021; Zhang *et al.*, 2020). Recently, it has been suggested that the high depletion of upper mantle elements is caused by melt dehydration above the 410 km discontinuity (Morikawa *et al.*, 2008; Stern, 2018; Umam *et al.*, 2022).

Geological Settings

Based on previous research studies, hot spring water in the Kinki region, southwestern Japan, is thought to have a close relationship with the subduction process (Umam *et al.*, 2026; Figure 1). One of the most well-known, the Arima hot spring water, has been confirmed to come from

deep groundwater as a result of the subduction process (Kusuda *et al.*, 2014; Nakamura *et al.*, 2016). However, along with the development of geochemical analysis, many researchers continue to conduct research to find analytical methods that are more sensitive and effective in analyzing geofluids. The development of geofluid analysis methods has become a special topic in the field of geoscience.

Over the past two decades, many researchers have analyzed several hot spring waters in the southwest of Japan and have speculated that some of the hot spring waters that appear are closely related to the subduction process by the Philippine Sea Slab (PHS). Various methods have been carried out, such as He gas analysis (Meibom *et al.*, 2003; Sano and Nakajima, 2008; Tedesco, 1997), tremor activity (Obara, 2002; Sano and Nakajima, 2008), geochemical analysis using REE (Nakamura *et al.*, 2014), and others to strengthen this suspicion.

Southwestern (SW) Japan is a nonvolcanic region that has many representative hot spring waters for research in recent years (Morikawa *et al.*, 2016). Some hot spring waters that appear




Figure 1. Schematic illustration of the tectonic setting of the Japanese Archipelago, showing the subduction of the Philippine Sea Plate beneath southwest Japan and its relationship to the occurrence of deep-seated fluids. The Arima hot spring area is highlighted, indicating the upward migration of slab-derived fluids generated during slab dehydration processes.

around the Median Tectonic Line (MTL) are in SW Japan (Morikawa *et al.*, 2016). MTL is a large fault that is still actively moving in the SW region of Japan, starting from the Shikoku region and extending to the Kinki region. Some researchers suspect that MTL movement activities have a contribution with hot spring water that occurs in the SW region of Japan (Taira, 2001; Tabei *et al.*, 2002; Iwamori, 2007). However, these approaches remain limited in resolving both fluid origin and geochemical processes simultaneously, particularly in nonvolcanic subduction settings. Therefore, an integrated geochemical framework is required.

This study aims to test whether a combined geochemical approach using lithium isotope systematics ($\delta^7\text{Li}$), stable isotopes (δD and $\delta^{18}\text{O}$), major and trace elements (*e.g.* Cl, Li, and Cl/Li ratios) can effectively detect and characterize the contribution of slab-dehydrated fluids to surface and shallow groundwater systems in the Arima region. To achieve this objective, previously published lithium isotope ($\delta^7\text{Li}$) data for four Arima samples reported by Umam *et al.* (2022) were re-evaluated together with Li and Cl concentration data reported by Umam *et al.* (2022) and Kusuda *et al.* (2014). The interpretation is based on comparing geochemical signatures, including $\delta^7\text{Li}$ values, Cl concentrations, and Cl/Li ratios, to distinguish between thermal water, meteoric water, and mixing processes.

The novelty of this study lies in the integrated use of lithium isotope systematics together with stable isotopes and major elements as a unified geochemical framework. While previous studies have primarily relied on single tracers (*e.g.* He isotopes or REE), this study proposes a multiparameter approach that may provide additional constraints on the identification and characterization of slab-derived fluids. Furthermore, this study explores the relationship between $\delta^7\text{Li}$ and Cl/Li ratios as indicators of fluid-rock interaction processes, including lithium adsorption and desorption mechanisms. This perspective provides new insights into the behaviour of lithium in hydrothermal systems and offers an alternative

interpretation beyond conventional dilution-based models. Overall, this integrated approach contributes to a more comprehensive understanding of slab-derived fluid signatures in non-volcanic subduction settings, particularly in the Arima hydrothermal system.

METHODS AND MATERIALS

Data Sources and Reanalysis

No new sampling, sample preparation, or geochemical measurements were conducted as part of the present study. Instead, this study integrated and re-evaluated previously published geochemical datasets for the Arima hydrothermal system. Lithium isotope ($\delta^7\text{Li}$) data for four Arima samples were obtained from Umam *et al.* (2022). Previously published Li and Cl concentration data were obtained from both Umam *et al.* (2022) and Kusuda *et al.* (2014) and were used to evaluate Li concentrations and calculate Cl/Li ratios. Stable isotope (δD and $\delta^{18}\text{O}$) and major-element data were also compiled from previously published studies, including Kusuda *et al.* (2014). Readers are referred to the respective original publications for information on the original samples and datasets.

The datasets from these previous studies were used according to the geochemical parameters available in each publication. Because the availability of individual parameters differs among the published datasets, the present study does not assume that all geochemical variables were measured for all Arima samples. Instead, each parameter was evaluated within the limits of the available published data and subsequently compared with the other geochemical indicators where appropriate. This approach allows the different published datasets to be considered within a common interpretative framework without implying that the original samples were collected or analyzed specifically for the present study.

Accordingly, $\delta^7\text{Li}$ was evaluated only for the samples for which lithium isotope data had previously been reported. For the broader Arima dataset,

Li and Cl concentrations and the corresponding Cl/Li relationships provide additional parameters that can be compared with the available stable-isotope and major-element information. The distinction between datasets with and without $\delta^7\text{Li}$ measurements is maintained throughout the present study so that the interpretation does not imply the existence of lithium isotope measurements where such data are unavailable.

Lithium isotope compositions are expressed in conventional delta notation as:

$$\delta^7\text{Li} (\text{‰}) = \left[\left(\frac{{}^7\text{Li}}{{}^6\text{Li}} \right)_{\text{sample}} / \left(\frac{{}^7\text{Li}}{{}^6\text{Li}} \right)_{\text{standard}} - 1 \right] \times 1000 \dots\dots\dots(1)$$

The $\delta^7\text{Li}$ values used in the present study are those previously reported by Umam *et al.* (2022); no new lithium isotope measurements or recalculation of the original isotope ratios were performed as part of the present study. Lithium isotope data were available only for the four Arima samples reported by Umam *et al.* (2022), whereas $\delta^7\text{Li}$ data were not available for the Arima samples reported by Kusuda *et al.* (2014).

In the present study, these previously published datasets were integrated and re-evaluated by comparing $\delta^7\text{Li}$ values, Li and Cl concentrations, calculated Cl/Li ratios, stable isotope compositions, and major-element chemistry. This integrated reanalysis was used to examine whether these geochemical parameters, individually and in combination, provide additional constraints on the characteristics and possible contribution of slab-dehydrated fluids to the Arima hydrothermal system.

This reanalysis therefore focuses on relationships among the published geochemical parameters rather than on reproducing the original analytical procedures. The previously reported values constitute the observational basis of the present study, whereas their integration and comparison within a common geochemical framework constitute the analytical approach applied here. In this way, the present study seeks to evaluate the extent to which multiple geochemical indicators can be used together to improve interpretation of fluid characteristics within the Arima hydrothermal system in Southwestern, Japan.

RESULT AND ANALYSIS

The previously published data for the four Arima samples reported by Umam *et al.* (2022) (Table 1) show high Chlorine (Cl) concentrations ranging from 15200 mg/L to 36500 mg/L. These Cl concentrations span values both below and above that of modern seawater (~19,000 mg/L) (Kazahaya *et al.*, 2014; Oi *et al.*, 1996), while remaining substantially higher than typical concentrations reported for shallow groundwater and surface water (Ii *et al.*, 2019).

Based on the data from Arima hot spring waters reported by Kusuda *et al.* (2014), sodium (Na) concentrations are high, ranging from 18 mg/L to 20,077 mg/L, and potassium (K) concentrations range from 3 mg/L to 2,593 mg/L. The concentration of calcium (Ca), magnesium (Mg), and sulphate (SO_4) range from 8 mg/L to 3,006 mg/L, 0 mg/L to 511 mg/L, and 0.7 mg/L to 59.02 mg/L.

Based on the Piper diagram (Figure 2), the Arima hot spring water samples reported by Kusuda *et al.* (2014) generally plot within the sodium chloride (Na-Cl) water type field. The sodium chloride type are mostly found in coastal area (Chandrasekar *et al.*, 2014; Satriani *et al.*, 2012), geothermal waters area (Deon *et al.*, 2015; Giggenbach, 1991; Millot *et al.*, 2012; Nukman and Hochstein, 2019; Siregar and Kurniawan, 2018; Giggenbach, 1992; Winters and Cawvey, 2015), or from the mantle (deep fluids) (Ii *et al.*, 2019; Kusuda *et al.*, 2014) in the groundwater category. Meanwhile, Piper diagrams are excellent for tracking the origins of early groundwater based on species dominance (Kumar *et al.*, 2009; Nazri *et al.*, 2016; Ravikumar and Somashekar, 2017; Singh *et al.*, 2020; Umar Kura *et al.*, 2013). However, further confirmation is still needed regarding the origin of the Arima hot water samples, whether they are derived from sea water intrusion, or underwater volcanoes, or from the mantle (deep fluids) due to the subduction process.

The interpretation results of the stable isotope (Figure 3) showed that Arima hot spring waters (Kusuda *et al.*, 2014) have a range of δD from -51.05 ‰ to -34.12 ‰ with a shift value $\delta^{18}\text{O}$ rang-

Table 1. Previously published Li and Cl concentrations and lithium isotope ($\delta^7\text{Li}$) data for Arima hot spring waters and seawater, together with calculated Cl/Li ratios

Fluid Type (Source)	Location	Sample Name	Li (ppm)	Cl/Li (wt./wt.)	Cl (ppm)	$\delta^7\text{Li}$ (‰)
Seawater (Oi <i>et al.</i> , 1996)	Sezaki, Japan	Seawater	0.17	111764.709	19000	30
Geofluids (Deep Groundwater Water) (Kusuda <i>et al.</i> , 2014)	Arima Type Fluids (Non Volcanic Area), Southwestern Japan	L1	38.16	436.740	16666	nm
		L2	40.75	526.110	21439	nm
		L3	6.5	521.846	3392	nm
		L4	0.03	466.666	14	nm
		L5	35.78	701.173	25088	nm
		L6	30.76	675.747	20786	nm
		L7	14.42	606.241	8742	nm
		L8	36.69	703.952	25828	nm
		L9	22.97	637.222	14637	nm
		L10	20.86	735.043	15333	nm
		L11	0.05	8700	435	nm
		L12	2.08	622.115	1294	nm
		L13	51.9	771.348	40033	nm
		L14	33.81	1082.579	36602	nm
Groundwater samples (Umam <i>et al.</i> , 2022)	Arima hot spring waters	A1	25.8	694	17900	1.5
		A2	46.4	733	34000	1.4
		A3	45.9	795	36500	1.5
		A4	27.2	559	15200	1.6

*nm: not measured




Figure 2. Piper diagram showing the hydrochemical facies of Arima hot spring waters (Kusuda *et al.*, 2014) compared with seawater (Oi *et al.*, 1996). Most samples plot within the Na-Cl type field, indicating a strong influence of deep-seated saline fluids. The clustering of samples near the sodium-chloride end member suggests a dominant contribution from high-salinity geothermal fluids rather than shallow meteoric sources. Minor deviations toward mixed-type fields indicate local mixing processes and secondary water-rock interaction.




Figure 3. Stable isotope composition (δD vs. $\delta^{18}\text{O}$) of Arima hot spring waters (Kusuda *et al.*, 2014) compared with seawater (Oi *et al.*, 1996), surface water (Hosono and Yamanaka, 2021), geothermal fluids from TVZ (Millot *et al.*, 2012), and magmatic water (Giggenbach, 1992). Most samples show a positive $\delta^{18}\text{O}$ shift from the Global Meteoric Water Line (GMWL), indicating significant water-rock interaction at elevated temperatures. The distribution of samples suggests mixing between deep geothermal fluids and meteoric water.

ing from -7.88 to 4.98 (from the Global Meteoric Water Line). According to Matsubaya *et al.* (1973), a high shift value occurs due to the long water-rock interaction process compared to surface or meteoric water. Furthermore, thermal water samples exhibit a higher shift than surface water and are similar to the geothermal waters from TVZ, New Zealand (Milot *et al.*, 2012). Based on the results of interpretation using stable isotopes, several Arima hot spring waters have plots that are close to other types of water, such as surface water and geothermal waters (Deon *et al.*, 2015; Giggenbach, 1991; Milot *et al.*, 2012; Nukman and Hochstein, 2019; Siregar and Kurniawan, 2018; Giggenbach, 1992; Winters and Cawvey, 2015). Thus, stable isotopes still require other evidence to more accurately detect the origin of Arima hot spring waters.

Previous studies have shown that Cl/Li ratios can provide useful information for characterizing deep-seated and hydrothermal fluids in Japan (Kazahaya *et al.*, 2014). In the present reanalysis, a Cl/Li ratio of 1,000 is used as a reference boundary for comparing variations among the previously published Arima datasets. Millot *et al.* (2012) and Tanimizu *et al.* (2021) confirmed that lithium (Li) is very sensitive to high temperatures. Furthermore, groundwater with a high Li concentration is commonly found in the inner layer, namely the mantle (Meredith *et al.*, 2013; Nishio *et al.*, 2010; Tang *et al.*, 2007; Tomascak, 2004). Information on the high Li concentration in groundwater supports the results of You *et al.* (1996) comparing the behaviour of Lithium (Li) and Boron (B) when experiencing temperature changes (Figure 4).




Figure 4. a). Experimental setup for simulating fluid-rock interaction under hydrothermal conditions (25-350°C, 800 bar) using an autoclave system with synthetic saline solution and sediment (You *et al.*, 1996); b). Temperature-dependent behaviour of Li and B during heating and cooling processes, highlighting the sensitivity of Li to thermal conditions and fluid evolution; c). A conceptual model linking slab dehydration processes to fluid migration, illustrating the generation of slab-derived fluids and their ascent toward the surface.

Based on the interpretation of Figure 5, the four Arima samples previously reported by Umam *et al.* (2022) have Li concentrations > 25 mg/L and Cl concentrations ranging from 15200

mg/L to 36500 mg/L. Meanwhile, the Arima hot spring water samples (Kusuda *et al.*, 2014) had Li concentrations ranging from 0.03 mg/L to 51.9 mg/L (Table 1), and several samples had Cl concentrations < 1,000 (Table 2).

The interpretation of Li concentrations and Cl/Li ratios (Figure 5) for the four Arima samples previously reported by Umam *et al.* (2022) shows Cl/Li values < 1000. Meanwhile, several samples of Arima hot spring waters (Kusuda *et al.*, 2014) has a Cl/Li value > 1,000. The results of research by Kazahaya *et al.* (2014) using forty-nine hot springs spread throughout Japan show that magmatic water has a Cl/Li > 1,000 (in weight ratio) with Cl > 1000 mg/L.

The results of the interpretation of lithium isotopes ($\delta^7\text{Li}$) and Cl/Li ratios (Figure 6) show that the four Arima samples previously reported by Umam *et al.* (2022) have $\delta^7\text{Li} < 5\text{‰}$ and Cl/Li values < 1000. Meanwhile, seawater (Oi *et al.*, 1996), which also has a Cl value > 1,000 mg/L, has $\delta^7\text{Li} \gg 5\text{‰}$. Based on the interpretation of Figure 6, $\delta^7\text{Li} = 5\text{‰}$ and Cl/Li = 1,000 are used as reference boundaries for comparing the geochemical characteristics of the available samples.




Figure 5. Relationship between Li concentration and Cl/Li ratios for groundwater samples (Umam *et al.*, 2022) and Arima hot spring waters (Kusuda *et al.*, 2014) compared with seawater (Oi *et al.*, 1996). The dashed line (Cl/Li = 1,000) is used as a reference boundary for comparing variations in Cl/Li ratios among the datasets. The four Arima samples previously reported by Umam *et al.* (2022) plot in the low Cl/Li (<1000) and high Li concentration field while several literature samples show higher Cl/Li values, indicating variability in fluid processes and possible mixing or geochemical modification.

Table 2. Previously published stable isotope and major-element data for Arima hot spring waters and seawater

Fluid Type (Source)	Location	Temp (°C)	Isotope stable		Major element (mg/L)						
			$\delta^{18}\text{O}$ (‰)	δD (‰)	Mg ²⁺	Na ⁺	K ⁺	Ca ²⁺	Cl ⁻	SO ₄	HCO ₃ ⁻
Seawater (Oi <i>et al.</i> , 1996)	Sezaki, Japan	12.6	0	0	1400	10600	431	934	19000	2470	nm
		22.3	-0.35	-40.44	284	9623	959	629	16666	6.54	3803
		25.9	1.09	-38.51	114	12618	1372	985	21439	0.7	3154
		32.1	-0.7	-51.05	35	1743	208	491	3392	1.12	781
		19.1	-7.88	-50.27	1	18	3	23	14	59.02	30
Geofluids (Deep Groundwater Water) (Kusuda <i>et al.</i> , 2014)	Arima Type Fluids (Non Volcanic Area), Southwestern Japan	95.3	2.07	-38.15	15	11945	2525	2192	25088	2.41	26
		88.3	0.77	-40.5	12	10127	2153	1845	20786	2.41	203
		80.3	-4.09	-45.92	15	4700	1000	627	8742	24.92	473
		88.3	2.22	-37.71	17	12189	2593	2356	25828	2.77	137
		61.5	-1.85	-44.69	29	7805	1388	1283	14637	10.3	435
		33.3	-3.26	-44.31	144	9014	332	860	15333	1.69	1623
		19.4	-7.27	-48.92	0	23	4	8	435	15.27	93
		19.5	-7.08	-48.56	4	704	131	108	1294	23.06	150
		29.2	4.98	-34.12	136	20077	1930	3006	40033	10.25	1293
		32.2	3.61	-35.11	511	18379	112	2598	36602	2.43	214

*nm: not measured




Figure 6. Relationship between lithium isotope composition ($\delta^7\text{Li}$) and Cl/Li ratios for groundwater samples (Umam *et al.*, 2022), compared with geothermal fluids from the Taupo Volcanic Zone (TVZ) (Millot *et al.*, 2012), and seawater (Oi *et al.*, 1996). Dashed lines at $\delta^7\text{Li} = 5\text{‰}$ and Cl/Li = 1,000 are shown as reference boundaries for comparing the distribution of the available samples rather than as definitive boundaries between three fluid categories. Groundwater samples cluster in the low $\delta^7\text{Li}$ (<5‰), and low Cl/Li (<1,000) field, consistent with deep hydrothermal fluids, while seawater plots in the high $\delta^7\text{Li}$ and high Cl/Li fields.

The four Arima samples reported by Umam *et al.* (2022) plot in the low- $\delta^7\text{Li}$ (<5‰)

and low-Cl/Li (<1,000) field, whereas seawater plots in the high- $\delta^7\text{Li}$ (>5‰) and high-Cl/Li (>1,000) field. Samples occurring in the other combinations of these two parameters should not be assigned to a specific fluid category solely on the basis of these boundaries and require consideration of additional geochemical evidence.

Even though the Arima data on hot spring waters (Kusuda *et al.*, 2014) does not have lithium isotope ($\delta^7\text{Li}$) data, based on interpretation of Li, Cl concentrations (Table 1) and Cl/Li ratios (Figure 5), as well as categories of fluid (Figure 6), almost all Arima hot spring water (Kusuda *et al.*, 2014) can be estimated to have Li > 0.1 mg/L, Cl > 1,000 mg/L, and Cl/Li < 1000, which are thermal water categories. On the other hand, one of the Arima hot spring waters (Kusuda *et al.*, 2014), which has a Li < 5 mg/L, Cl < 1,000 mg/L, and Cl/Li > 1,000, is not in the thermal water category, but it is likely to be in the mixing water category because it has a fairly high Cl value, namely, 435 mg/L (Table 1), and the sample collection location is in the Arima hot spring water area (far from the coastal area).

DISCUSSION

The combined interpretation of major elements (Figure 2), stable isotopes (Figure 3), and lithium systematics (Figure 5 and Figure 6) provides a progressively refined understanding of the origin and evolution of Arima hydrothermal fluids. The Piper diagram (Figure 2) shows that most samples belong to the Na-Cl type, indicating a strong influence of saline, deep-seated fluids (Kusuda *et al.*, 2014). However, this hydrochemical signature is not unique and may also result from seawater intrusion or prolonged water-rock interaction. Similarly, stable isotope data (Figure 3) exhibit a clear $\delta^{18}\text{O}$ shift indicative of high-temperature water-rock interaction, but overlapping fields between meteoric, geothermal, and mixed waters limit their ability to uniquely constrain fluid origin.

Additional ambiguity is observed in the Cl/Li relationships (Figure 5), where inconsistent patterns occur within the same hydrothermal system. Some samples display Cl/Li ratios traditionally associated with deep fluids, whereas others show contrasting behaviour that can not be explained solely by mixing or dilution. This inconsistency highlights a key limitation of using Cl/Li as a direct tracer of fluid sources. The integration of lithium isotope systematics (Figure 6) resolves this ambiguity. The groundwater samples consistently plot in the low $\delta^7\text{Li}$ (<5‰) and low Cl/Li (<1,000) fields, clearly distinguishing them from seawater and indicating a strong contribution from deep hydrothermal or slab-derived fluids. Unlike major elements and stable isotopes, $\delta^7\text{Li}$ provides a more direct and process-sensitive constraint on fluid origin due to its response to water-rock interaction (Umam *et al.*, 2024).

Importantly, the combined $\delta^7\text{Li}$ -Cl/Li framework demonstrates that Cl/Li ratios primarily reflect lithium mobility controlled by adsorption-desorption processes rather than fluid sources. This reinterpretation explains the observed inconsistencies and indicates that variations in Cl/Li are governed by different stages of fluid-rock interaction. The integration of Li concentration and Cl/Li-based hydrochemical classification into the

stable isotope framework (Figure 7) provides additional constraints on fluid interpretation. Although δD - $\delta^{18}O$ data alone show overlapping fields, the addition of Li concentration and Cl/Li-based hydrochemical classification provides an additional basis for distinguishing thermal water from mixing water. Thermal waters exhibit larger $\delta^{18}O$ shifts, reflecting stronger water-rock interaction, whereas mixing waters plot closer to the meteoric field due to dilution effects (Umam *et al.*, 2026).

Samples with similar stable isotope signatures can belong to different fluid categories when Li concentrations and Cl/Li relationships are considered. This demonstrates that stable isotopes alone may mask differences in fluid origin and evolution. Overall, this study shows that conventional tracers (major elements and stable isotopes) are insufficient to resolve fluid origin in complex hydrothermal systems (Siregar *et al.*, 2026a; Siregar *et al.*, 2026b). The integration of lithium

isotope systematics may provide an additional process-sensitive constraint that improves the interpretation of possible slab-dehydrated fluid contributions. This combined approach establishes a new geochemical framework in which:

- δ^7Li serves as a tracer of fluid origin and water-rock interaction.
- Cl/Li reflects lithium mobility controlled by adsorption-desorption processes.

Such a framework may provide additional constraints for evaluating possible slab-derived fluid contributions in non-volcanic subduction settings, including the Arima Hydrothermal System.

By integrating Li and Cl concentrations and Cl/Li-based hydrochemical classification with stable isotope and major element, additional constraints on fluid types and characteristics can be obtained (Figure 7 and Figure 8). This combined approach enhances the resolution of fluid classification beyond what can be obtained from individual tracers. Figure 7 shows




Figure 7. Stable isotope composition (δD - $\delta^{18}O$) of Arima hot spring waters combined with lithium-based hydrochemical classification. The diagram shows δD versus $\delta^{18}O$ values of Arima hot spring waters, together with reference fields for seawater, magmatic water, geothermal waters, surface water, and mixing water. Based on the hydrochemical criteria of Kusuda *et al.* (2014), most Arima hot spring waters are classified as thermal waters, characterized by relatively high Li concentrations (>0.1 mg/L), high Cl concentrations ($>1,000$ mg/L), and low Cl/Li ratios ($<1,000$) (Table 1). In contrast, one sample with lower Cl concentration, lower Li concentration, and higher Cl/Li ratio is interpreted as mixing water (see Figure 5). Although lithium isotope data are not plotted in this figure, Li concentration and Cl/Li ratios are used to support the hydrochemical classification of the plotted samples.




Figure 8. Piper diagram integrated with Li concentration and Cl/Li-based hydrochemical classification for Arima hot spring waters. Red circles represent thermal water, while green circles indicate mixing water, based on Li and Cl concentrations and Cl/Li relationships. Most samples cluster within the Na-Cl type field, while mixing water samples show slight deviations, reflecting dilution and interaction with shallow groundwater. This integrated approach suggests that similar hydrochemical facies may represent different fluid characteristics when Li concentrations and Cl/Li relationships are considered together with major-element compositions.

that several Arima hot spring waters (Kusuda *et al.*, 2014) exhibit relatively small $\delta^{18}\text{O}$ shifts and plot near the surface water field. However, when Li concentrations and Cl/Li relationships are considered together with the stable isotope data, these samples may be further distinguished according to their fluid characteristics. Samples represented by red circles correspond to thermal waters dominated by slab-derived fluids, while green circles indicate mixing between thermal water and surface or shallow groundwater.

This classification is consistent with the major element interpretation (Figure 8), where the mixing water samples (green circles) still exhibit relatively high Cl concentrations (*e.g.* ~435 mg/L; Table 1) and fall within the Na-Cl type field. This suggests that even mixed waters retain a significant contribution from deep hydrothermal fluids. Although many samples from Kusuda *et al.* (2014) plot within the same Na-Cl field, the integration of Li concentrations and Cl/Li-based hydrochemical classification may provide additional discrimination of fluid characteristics and evolutionary processes. This demonstrates that similar hydrochemical facies can mask fundamentally different fluid histories.

Most samples classified as thermal water (red circles) cluster within the Na-Cl type field, confirming the dominant influence of deep, saline hydrothermal fluids. This is consistent with previous interpretations based on major element chemistry. However, the incorporation of Li concentrations and Cl/Li relationships suggests that samples within the same Na-Cl facies may exhibit different geochemical characteristics or evolutionary histories. Mixing water samples (green circles) also plot near the Na-Cl field but show slight deviations toward other hydrochemical facies, reflecting dilution by surface or shallow groundwater. Despite these deviations, their relatively high Cl concentrations indicate that they still retain a significant contribution from deep hydrothermal fluids (Umam *et al.*, 2025).

Importantly, Figure 8 demonstrates that hydrochemical classification alone is insufficient to distinguish fluid origin, as both thermal and

mixing waters can occupy similar positions in the Piper diagram. The addition of Li concentration and Cl/Li-based hydrochemical classification may help distinguish these fluid types, even when their major-element compositions appear similar. This result reinforces the interpretation that major element chemistry primarily reflects bulk composition, while lithium isotopes provide insight into fluid origin and geochemical processes. The combined approach therefore offers a more comprehensive framework for interpreting hydrothermal systems.

CONCLUSIONS

This study suggests that the integration of lithium isotope systematics, stable isotopes, and major element geochemistry may provide a useful framework for evaluating the origin and evolution of hydrothermal fluids in the Arima System, southwestern Japan. The previously published Li concentration and lithium isotope ($\delta^7\text{Li}$) data for the four Arima samples reported by Umam *et al.* (2022) show low $\delta^7\text{Li}$ values (<5‰) and low Cl/Li ratios (<1000), clearly distinguishing them from seawater, which exhibits significantly higher $\delta^7\text{Li}$ values and Cl/Li ratios. Cl/Li, three fluid categories can be defined: (1) thermal water ($\delta^7\text{Li}$ <5 ‰ and Cl/Li <1,000), (2) mixing water between thermal and surface water, and (3) surface or shallow groundwater ($\delta^7\text{Li}$ >5 ‰ and Cl/Li >1,000). The interpretation of these previously published Arima data in the present study places the four samples within the thermal water category, indicating a strong contribution from slab-dehydrated fluids.

Although lithium isotope data are not available for the Arima hot spring waters reported in previous studies (Kusuda *et al.*, 2014), their Li concentrations (>0.1 mg/L), Cl concentrations (>1,000 mg/L), and Cl/Li ratios (<1,000) suggest that most samples also belong to the thermal water category. In contrast, one sample with relatively low Li (<5 mg/L), low Cl (<1000 mg/L), and high Cl/Li (>1000), despite having moderate Cl (~435

mg/L), is interpreted as mixing water, reflecting interaction between thermal fluids and surface or shallow groundwater.

Major element compositions (Piper diagram) indicate that most samples belong to the Na-Cl type, suggesting a strong influence of saline deep fluids. However, this signature alone is not diagnostic of fluid origin. Similarly, stable isotopes (δD - $\delta^{18}O$) indicate significant water-rock interaction but cannot uniquely distinguish between fluid sources due to overlapping fields. Furthermore, this study suggests that Cl/Li ratios may be influenced by lithium mobility associated with adsorption-desorption processes as well as by fluid-source characteristics, which may help explain some of the variability observed when Cl/Li is used as a standalone tracer.

In contrast, lithium isotope systematics (δ^7Li) provide a more sensitive and process-based tracer, enabling clearer discrimination between thermal, mixed, and surface waters. The integration of lithium isotopes into stable isotope and major element interpretations may provide additional constraints on fluid classification and help reduce ambiguities associated with individual geochemical tracers.

Overall, this study proposes an integrated geochemical framework in which δ^7Li may provide information on fluid origin and water rock interaction, while Cl/Li may reflect the combined influence of lithium mobility, fluid-rock interaction, and fluid characteristics. This combined approach may provide additional constraints for evaluating possible slab-dehydrated fluid contributions in nonvolcanic subduction settings and may also be applicable to other hydrothermal systems.

ACKNOWLEDGEMENTS

The authors would like to express sincere gratitude to Professor Masaharu Tanimizu, Dr Yasunori Mori, Mr Ryo Nakai, and Mr Naoto Sugimoto for their valuable support and assistance during the hot spring water sampling in The Kii Peninsula,

Japan. Their expertise, guidance, and generous collaboration greatly contributed to the success of the data collection process. The author would like to thank anonymous reviewers for their constructive reviews. This research was funded by The Nishimura Foundation from Mandom Company, Japan, and The Centre for Research in Radiation, Isotopes, and Earth System Sciences (CRiES), University of Tsukuba, Japan.

Conflicts of Interest

The authors declare no conflict of interest concerning the publication of this article. The authors also confirm that the data and the article are free of plagiarism.

REFERENCES

- Chandrasekar, N., Selvakumar, S., Srinivas, Y., John Wilson, J.S., Simon Peter, T., and Magesh, N.S., 2014. Hydrogeochemical assessment of groundwater quality along the coastal aquifers of southern Tamil Nadu, India. *Environmental Earth Sciences*, 71 (11), p.4739-4750. <https://doi.org/10.1007/>.
- Deon, F., Förster, H.J., Brehme, M., Wiegand, B., Scheytt, T., Moeck, I., Jaya, M.S., and Putriatni, D. J., 2015 . Geochemical/hydrochemical evaluation of the geothermal potential of the Lamongan volcanic field (Eastern Java, Indonesia). *Geothermal Energy*, 3 (1), p.1-21. <https://doi.org/10.1186>.
- Giggenbach, W.F., 1991. Chemical Techniques in Geothermal Exploration. In *Chemistry Division*, p.119-144.
- Giggenbach, W.F., 1992. Isotopic shifts in waters from geothermal and volcanic systems along convergent plate boundaries and their origin. *Earth and Planetary Science Letters*, 113 (4), p.495-510.
- Giggenbach, W. F., 1992. Isotopic shifts in waters from geothermal and volcanic systems along convergent plate boundaries and their origin. *Earth and Planetary Science Letters*, 113 (4), p.495-510.

- Harlaux, M., Mercadier, J., Bonzi, W.M.E., Kremer, V., Marignac, C., and Cuney, M., 2017. Geochemical Signature of Magmatic-Hydrothermal Fluids Exsolved from the Beauvoir Rare-Metal Granite (Massif Central, France): Insights from LA-ICPMS Analysis of Primary Fluid Inclusions. *Geofluids*, 2017, 25pp. <https://doi.org/10.1155/2017/1925817>.
- Hosono, T. and Yamanaka, C., 2021. Origins and pathways of deeply derived carbon and fluids observed in hot spring waters from non-active volcanic fields, western. *Earth, Planets and Space*, 73:155, p.1-12. <https://doi.org/10.1186/s40623-021-01478-1>.
- Ii, H., Kitagawa, H., Kubohara, T., and Machida, I., 2019. Characteristics of water chemistry for Arima-type deep thermal water in the Kinokawa River catchment, Kii Peninsula, Japan. *International Journal of GEOMATE*, 17 (62), p.158-166. <https://doi.org/10.21660/2019.62.7156>.
- Iwamori, H., 2007. Transportation of H₂O beneath the Japan arcs and its implications for global water circulation. *Chemical Geology*, 239 (3-4), p.182-198. <https://doi.org/10.1016/j.chemgeo.2006.08.011>.
- Kazahaya, K., Takahashi, M., Yasuhara, M., Nishio, Y., Inamura, A., Morikawa, N., Sato, T., Takahashi, H.A., Kitaoka, K., Ohsawa, S., Oyama, Y., Ohwada, M., Tsukamoto, H., Horiguchi, K., Tosaki, Y., and Kirita, T., 2014. *Spatial distribution and feature of slab-related deep-seated fluid in SW Japan*, 44 (1), p.3-16. <https://doi.org/10.4145/jahs.44.3>.
- Kumar, S.K., Rammohan, V., Sahayam, J.D., and Jeevanandam, M., 2009. Assessment of groundwater quality and hydrogeochemistry of Manimuktha River basin, Tamil Nadu, India. *Environmental Monitoring and Assessment*, 159 (1-4), p.341-351. <https://doi.org/10.1007/s10661-008-0633-7>.
- Kusuda, C., Iwamori, H., Nakamura, H., Kazahaya, K., and Morikawa, N., 2014. Arima hot spring waters as a deep-seated brine from subducting slab. *Earth, Planets and Space*, 66:119, p.1-13. <https://doi.org/10.1186/1880-5981-66-119>.
- Matsubaya, O., Sakai, H., Kusachi, I., and Satake, H., 1973. Hydrogen and oxygen isotopic ratios and major element chemistry of Japanese thermal water systems. *Geochemical Journal*, 7 (3), p.123-151. <https://doi.org/10.2343/geochemj.7.123>.
- Meibom, A., Anderson, D.L., Sleep, N.H., Frei, R., Chamberlain, C.P., Hren, M.T., and Wooden, J.L., 2003. Are high ³He/⁴He ratios in oceanic basalts an indicator of deep-mantle plume components? *Earth and Planetary Science Letters*, 208 (3-4), p.197-204. [https://doi.org/10.1016/S0012-821X\(03\)00038-4](https://doi.org/10.1016/S0012-821X(03)00038-4).
- Meredith, K., Moriguti, T., Tomascak, P., Hollins, S., and Nakamura, E., 2013. The lithium, boron and strontium isotopic systematics of groundwaters from an arid aquifer system: Implications for recharge and weathering processes. *Geochimica et Cosmochimica Acta*, 112, p.20-31. <https://doi.org/10.1016/j.gca.2013.02.022>.
- Millot, R., Hegan, A., and Negrel, P., 2012. Geothermal waters from the Taupo Volcanic Zone, New Zealand: Li, B, and Sr isotopes characterization. *Applied Geochemistry*, 27, p.677-688. <https://doi.org/doi:10.1016/j.apgeochem.2011.12.015>.
- Morikawa, N., Kazahaya, K., Masuda, H., Ohwada, M., Nakama, A., Nagao, K., and Sumino, H., 2008. Relationship between geological structure and helium isotopes in deep groundwater from the Osaka Basin: Application to deep groundwater hydrology. *Geochemical Journal*, 42 (1), p.61-74. <https://doi.org/10.2343/geochemj.42.61>.
- Morikawa, N., Kazahaya, K., Takahashi, M., Inamura, A., Takahashi, H. A., Yasuhara, M., Ohwada, M., Sato, T., Nakama, A., Handa, H., Sumino, H., and Nagao, K., 2016. Widespread distribution of ascending fluids transporting mantle helium in the fore-arc region and their upwelling processes: Noble gas and major element composition of deep groundwater in the Kii Peninsula, southwest Japan. *Geochimica et Cosmochimica Acta*, 182, p.173-196. <https://doi.org/10.1016/j.gca.2016.03.017>.

- Nakajima, J. and Hasegawa, A., 2007. Subduction of the Philippine Sea plate beneath southwestern Japan: Slab geometry and its relationship to arc magmatism. *Journal of Geophysical Research: Solid Earth*, 112 (8), p.1-18. <https://doi.org/10.1029/2006JB004770>.
- Nakamura, H., Fujita, Y., Nakai, S., Yokoyama, T., and Iwamori, H., 2014. Rare Earth Elements and Sr-Nd-Pb isotopic analyses of the Arima hot spring waters, Southwest Japan: Implications for the origin of the Arima-type brine. *Journal of Geology and Geosciences*, 3 (4), p.1-8. <https://doi.org/10.4172/2329-6755.1000161>
- Nakamura, H., Chiba, K., Chang, Q., Morikawa, N., Kazahaya, K., and Iwamori, H., 2016. Origin of the Arima-type and Associated Spring Waters in the Kinki District, Southwest Japan. *Journal of Geology and Geophysics*, 05:02, p.1-15. <https://doi.org/10.4172/2381-8719.1000240>.
- Nazri, M.A.A., Tan, L.W., Kasmin, H., Syafalni, S., and Abustan, I., 2016. Geophysical and Hydrochemical Characteristics of Groundwater at Kerian Irrigation Scheme. *IOP Conference Series: Materials Science and Engineering*, 136, 12070. <https://doi.org/10.1088/1757-899X/136/1/012070>.
- Nishio, Y., Okamura, K., Tanimizu, M., Ishikawa, T., and Sano, Y., 2010. Lithium and strontium isotopic systematics of waters around Ontake volcano, Japan: Implications for deep-seated fluids and earthquake swarms. *Earth and Planetary Science Letters*, 297 (3-4), p.567-576. <https://doi.org/10.1016/j.epsl.2010.07.008>.
- Nukman, M. and Hochstein, M.P., 2019. The Sipoholon Geothermal Field and adjacent geothermal systems along the North-Central Sumatra Fault Belt, Indonesia: Reviews on geochemistry, tectonics, and natural heat loss. *Journal of Asian Earth Sciences*, 170 (November 2018), p.316-328. <https://doi.org/10.1016/j.jseaes.2018.11.007>.
- Obara, K., 2002. Nonvolcanic deep tremor associated with subduction in southwest Japan. *Science*, 296 (5573), p.1679-1681. <https://doi.org/10.1126/science.1070378>.
- Oi, T., Ikeda, K., Nakano, M., Ossaka, T., and Ossaka, J., 1996. Boron isotope geochemistry of hot spring waters in Ibusuki and adjacent areas, Kagoshima, Japan. *Geochemical Journal*, 30 (5), p.273-287. <https://doi.org/10.2343/geochemj.30.273>.
- Ravikumar, P. and Somashekar, R.K., 2017. Principal component analysis and hydrochemical facies characterization to evaluate groundwater quality in Varahi river basin, Karnataka state, India. *Applied Water Science*, 7 (2), p.745-755. <https://doi.org/10.1007/s13201-015-0287-x>.
- Sano, Y. and Nakajima, J., 2008. Geographical distribution of $^3\text{He}/^4\text{He}$ ratios and seismic tomography in Japan. *Geochemical Journal*, 42 (1), p.51-60. <https://doi.org/10.2343/geochemj.42.51>.
- Satriani, A., Loperte, A., Imbrenda, V., and Lappenna, V., 2012. Geoelectrical surveys for characterization of the coastal saltwater intrusion in metapontum forest reserve (Southern Italy). *International Journal of Geophysics*, 2012. <https://doi.org/10.1155/2012/238478>.
- Shiomi, K. and Park, J., 2008. Structural features of the subducting slab beneath the Kii Peninsula, central Japan: Seismic evidence of slab segmentation, dehydration, and anisotropy. *Journal of Geophysical Research: Solid Earth*, 113 (10), p.1-13. <https://doi.org/10.1029/2007JB005535>.
- Singh, K.K., Tewari, G., and Kumar, S., 2020. Evaluation of Groundwater Quality for Suitability of Irrigation Purposes: A Case Study in the Udham Singh Nagar, Uttarakhand. *Journal of Chemistry*, 2020. <https://doi.org/10.1155/2020/6924026>.
- Siregar, R.N. and Kurniawan, W.B., 2018. 2D Interpretation of Subsurface Hot Spring Geothermal Structure In Nyelanding Village Through Schlumberger Geoelectricity. *Jurnal Ilmiah Pendidikan Fisika Al-BiRuNi*, 07 (April), p.81-87. <https://doi.org/10.24042/jipfalbiruni.v7i1.2324>.

- Siregar, R.N., Sismanto, S., Triyana, K., Harijoko, A., Iqbal, M., Antarnusa, G., Putra, F.G., and Umam, R., 2026a. Lithium enrichment in high radiogenic geothermal systems originating from lithospheric water due to water-rock interactions. *Geothermics*, 134, 103499. <https://doi.org/10.1016/j.geothermics.2025.103499>.
- Siregar, R.N., Umam, R., Harijoko, A., Iqbal, M., Puspitarum, D.L., and Sismanto, S., 2026b. Radiogenic geothermal fluid from Bangka Island, Indonesia: Genesis and reservoir temperature estimation. *Geosystems and Geoenvironment*, 3 (3), 100537. <https://doi.org/10.1016/j.geogeo.2026.100537>.
- Stern, R.J., 2018. The evolution of plate tectonics. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 376 (2132). <https://doi.org/10.1098/rsta.2017.0406>.
- Tabei, T., Hashimoto, M., Miyazaki, S., Hirahara, K., Kimata, F., Matsushima, T., Tanaka, T., Eguchi, Y., Takaya, T., Hosono, Y., Ohya, F., and Kato, T., 2002. Subsurface structure and faulting of the Median Tectonic Line, southwest Japan inferred from GPS velocity field. *Earth, Planets and Space*, 54 (11), p.1065-1070. <https://doi.org/10.1186/BF03353303>.
- Taira, A., 2001. Tectonic Evolution of the Japanese Island Arc System. *Ocean Research*, 180 (September), p.1-2.
- Tang, Y.J., Zhang, H.F., and Ying, J.F., 2007. Review of the lithium isotope system as a geochemical tracer. *International Geology Review*, 49 (4), p.374-388. <https://doi.org/10.2747/0020-6814.49.4.374>.
- Tanimizu, M., Sugimoto, N., Hosono, T., Kuribayashi, C., Morimoto, T., Ito, A., and Umam, R., 2021. Application of B and Li isotope systematics for detecting chemical disturbance in groundwater associated with large shallow inland earthquakes in Kumamoto, Japan. 55, p.241-250. <https://doi.org/10.2343/geochemj.2.0633>.
- Tedesco, D., 1997. Systematic variations in the $^3\text{He}/^4\text{He}$ ratio and carbon of fumarolic fluids from active volcanic areas in Italy: Evidence for radiogenic ^4He and crustal carbon addition by the subducting African plate? *Earth and Planetary Science Letters*, 151 (3-4), p.255-269. [https://doi.org/10.1016/s0012-821x\(97\)81852-3](https://doi.org/10.1016/s0012-821x(97)81852-3).
- Tomaschak, P.B., 2004. Developments in the understanding and application of lithium isotopes in the earth and planetary sciences. *Reviews in Mineralogy and Geochemistry*, 55 (1), p.153-195. <https://doi.org/doi.org/10.2138/gsrmg.55.1.153>.
- Umam, R., Tanimizu, M., Nakamura, H., Nishio, Y., Nakai, R., Sugimoto, N., Mori, Y., Kobayashi, Y., Ito, A., Wakaki, S., Nagaishi, K., and Ishikawa, T., 2022. Lithium isotope systematics of Arima hot spring waters and groundwaters in Kii Peninsula. *Geochemical Journal*, 56 (5), p.8-17. <https://doi.org/10.2343/geochemj.GJ22015>.
- Umam, R., Cengiz, K., and Said, A., 2024. Application of major and trace elements for detecting the origin of groundwater: Lithium enrichment in Ain Al-Harrah hot spring influenced by the Red Sea, Saudi Arabia. *International Journal of Hydrology and Environmental Sustainability*, 3 (3), p.151-162. <https://doi.org/10.58524/ijhes.v3i3.522>.
- Umam, R., Junaidi, R., Syazali, M., Farid, F., Saregar, A., and Andiyani, A., 2025. Optimization of Piper trilinear diagram using lithium isotope systematics: An application for detecting the contribution of geothermal water from Aso Caldera after the 2016 earthquake in the Kumamoto Aquifer, Japan. *Indonesian Journal of Science and Technology*, 10 (1), p.159-170. <https://doi.org/10.17509/ijost.v10i1.80974>.
- Umam, R., Adachi, I., and Yamanaka, T., 2026. H–O–B–Li isotope systematics for assessing slab-derived fluids: Importance of post-dehydration processes. *Applied Geochemistry*, 197, 106667. <https://doi.org/10.1016/j.apgeochem.2025.106667>.
- Umar Kura, N., Firuz Ramli, M., Azmin Sulaiman, W. N., Ibrahim, S., Zaharin Aris, A., and Mustapha, A., 2013. Evaluation of factors

- influencing the groundwater chemistry in a small tropical Island of Malaysia. *International Journal of Environmental Research and Public Health*, 10 (5), p.1861-1881. <https://doi.org/10.3390/ijerph10051861>.
- Winters, M.S. and Cawvey, M., 2015. Governance obstacles to geothermal energy development in Indonesia. *Journal of Current Southeast Asian Affairs*, 34 (1), p.27-56. <https://doi.org/10.1177/186810341503400102>.
- Wyllie, P.J., 1974. Plate Tectonics, Seafloor Spreading, and Continental Drift: An Introduction. In *Plate Tectonics Assessments and Reassessments* (Charles F, Issue 1 858, p. 23).
- You, C.F., Castillo, P.R., Gieskes, J.M., Chan, L H., and Spivack, A.J., 1996. Trace element behaviour in hydrothermal experiments: Implications for fluid processes at shallow depths in subduction zones. *Earth and Planetary Science Letters*, 140 (1-4), p.41-52. [https://doi.org/10.1016/0012-821X\(96\)00049-0](https://doi.org/10.1016/0012-821X(96)00049-0).
- Zhang, K., Peng, J., Liu, W., Li, B., Xia, Q., Cheng, S., Yang, Y., Zeng, Y., Wen, M., Liu, D., Huang, Y., Zhong, L., Liu, P., and Jiang, X., 2020. The role of deep geofluids in the enrichment of sedimentary organic matter: A case study of the late ordovician-early silurian in the upper yangtze region and early cambrian in the lower Yangtze region, South China. *Geofluids*, 2020. <https://doi.org/10.1155/2020/8868638>.